


[CDC Home](#) | [Search](#) | [Health Topics A-Z](#)

National Center for Chronic Disease Prevention &amp; Health Promotion

**Behavioral Risk Factor Surveillance System**[BRFSS Home](#) | [Contact Us](#)
[Select Another Question](#)
[Remove Comparison](#)
[List All States](#)
**Prevalence Data**
**2004 - Washington vs  
California  
Alcohol Consumption**

 Grouped by Age 
**Binge drinkers (adults having five or more drinks on one occasion)**

To view a different grouping, choose from the grouping list and click the 'Go' button. To see all groupings for a response, click the response link in the table.

 Age:  

State:		Yes	No
Washington	% CI n	28.8 (25.6-31.9) 360	71.1 (68.0-74.3) 983
California	% CI n	22.0 (16.8-27.2) 77	77.9 (72.7-83.1) 241

% = Percentage, CI = Confidence Interval, n = Cell Size


Percentages are weighted to population characteristics.

Use caution in interpreting cell sizes less than 50.

N/A = Not available if the unweighted sample size for the denominator was &lt; 50 or the CI half width was &gt; 10 for any cell, or if the state did not collect data for that calendar year.

 See [Notes for data users](#) and [States conducting surveillance](#), by year.


**5 or more drinks on 1 occasion  
Washington vs California - 2004  
Age: 18-24**



Compare Washington - 2004 to Another Year:

2004

Compare

Compare Washington - 2004 to Another State:

Washington

Compare

[Privacy Policy](#) | [Accessibility](#)[BRFSS Home](#) | [Contact Us](#)[CDC Home](#) | [Search](#) | [Health Topics A-Z](#)

This page last reviewed June 15, 2005

[United States Department of Health and Human Services](#)  
[Centers for Disease Control and Prevention](#)  
[National Center Chronic Disease Prevention and Health Promotion](#)  
[Division of Adult and Community Health](#)


[CDC Home](#)
[Search](#)
[Health Topics A-Z](#)

National Center for Chronic Disease Prevention &amp; Health Promotion

**Behavioral Risk Factor Surveillance System**
[BRFSS Home](#) | [Contact Us](#)
[Select Another Question](#)
[Remove Comparison](#)
[List All States](#)
**Prevalence Data**
**2004 - Washington vs  
Nationwide  
Alcohol Consumption**

Grouped by Age

**Binge drinkers (adults having five or more drinks on one occasion)**

To view a different grouping, choose from the grouping list and click the 'Go' button. To see all groupings for a response, click the response link in the table.

Age: 18-24



State:		Yes	No
Washington	% CI n	28.8 (25.6-31.9) 360	71.1 (68.0-74.3) 983
Nationwide	Median % # States	28.1 52	71.8 52

% = Percentage, CI = Confidence Interval, n = Cell Size

Percentages are weighted to population characteristics.

Use caution in interpreting cell sizes less than 50.

N/A = Not available if the unweighted sample size for the denominator was < 50 or the CI half width was > 10 for any cell, or if the state did not collect data for that calendar year.


# of States Includes District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands in applicable years.

See [Notes for data users](#) and [States conducting surveillance](#), by year.

[http://apps.nccd.cdc.gov/brfss/age\\_c.asp?ctype=C&qtype\\_c=C&bkey=20045309&yr=2004&state=WA&q...](http://apps.nccd.cdc.gov/brfss/age_c.asp?ctype=C&qtype_c=C&bkey=20045309&yr=2004&state=WA&q...) 3/6/2006

TX315\_003

**5 or more drinks on 1 occasion  
Washington vs Nationwide - 2004  
Age: 18-24**



Compare Washington - 2004 to Another Year:

2004

Compare

Compare Washington - 2004 to Another State:

Washington

Compare

[Privacy Policy](#) | [Accessibility](#)[BRFSS Home](#) | [Contact Us](#)[CDC Home](#) | [Search](#) | [Health Topics A-Z](#)

This page last reviewed June 15, 2005

[United States Department of Health and Human Services](#)  
[Centers for Disease Control and Prevention](#)  
[National Center Chronic Disease Prevention and Health Promotion](#)  
[Division of Adult and Community Health](#)




[CDC Home](#)
[Search](#)
[Health Topics A-Z](#)

National Center for Chronic Disease Prevention &amp; Health Promotion

**Behavioral Risk Factor Surveillance System**
[BRFSS Home](#) | [Contact Us](#)
[Select Another Question](#)
[Remove Comparison](#)
[List All States](#)
**Prevalence Data**
**2004 - Washington vs  
California  
Alcohol Consumption**

Grouped by Age

**Adults who have had at least one drink of alcohol within the past 30 days**

To view a different grouping, choose from the grouping list and click the 'Go' button. To see all groupings for a response, click the response link in the table.

Age: 18-24



State:		Yes	No
Washington	% CI n	60.8 (57.5-64.0) 791	39.1 (35.9-42.4) 558
California	% CI n	54.5 (47.9-61.1) 183	45.4 (38.8-52.0) 136

% = Percentage, CI = Confidence Interval, n = Cell Size

Percentages are weighted to population characteristics.

Use caution in interpreting cell sizes less than 50.


N/A = Not available if the unweighted sample size for the denominator was < 50 or the CI half width was > 10 for any cell, or if the state did not collect data for that calendar year.

See [Notes for data users](#) and [States conducting surveillance, by year](#).

[http://apps.nccd.cdc.gov/brfss/age\\_c.asp?qtype=C&qtype\\_c=C&bkey=20045308&yr=2004&state=WA&q...](http://apps.nccd.cdc.gov/brfss/age_c.asp?qtype=C&qtype_c=C&bkey=20045308&yr=2004&state=WA&q...) 3/6/2006

TX315\_005

## At least one drink of alcohol in past 30 days Washington vs California - 2004 Age: 18-24



Compare Washington - 2004 to Another Year:

2004

Compare

Compare Washington - 2004 to Another State:

Washington

Compare

[Privacy Policy](#) | [Accessibility](#)[BRFSS Home](#) | [Contact Us](#)[CDC Home](#) | [Search](#) | [Health Topics A-Z](#)

This page last reviewed June 15, 2005

[United States Department of Health and Human Services](#)  
[Centers for Disease Control and Prevention](#)  
[National Center Chronic Disease Prevention and Health Promotion](#)  
[Division of Adult and Community Health](#)


[CDC Home](#)
[Search](#)
[Health Topics A-Z](#)

National Center for Chronic Disease Prevention &amp; Health Promotion

**Behavioral Risk Factor Surveillance System**
[BRFSS Home](#) | [Contact Us](#)
[Select Another Question](#)
[Remove Comparison](#)
[List All States](#)
**Prevalence Data**
**2004 - Washington vs  
Nationwide  
Alcohol Consumption**

Grouped by Age

Go

**Adults who have had at least one drink of alcohol within the past 30 days**

To view a different grouping, choose from the grouping list and click the 'Go' button. To see all groupings for a response, click the response link in the table.

Age: 18-24

Go

State:		Yes	No
Washington	% CI n	60.8 (57.5-64.0) 791	39.1 (35.9-42.4) 558
Nationwide	Median % # States	58.4 52	40.8 52

% = Percentage, CI = Confidence Interval, n = Cell Size

Percentages are weighted to population characteristics.

Use caution in interpreting cell sizes less than 50.

N/A = Not available if the unweighted sample size for the denominator was < 50 or the CI half width was > 10 for any cell, or if the state did not collect data for that calendar year.


# of States includes District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands in applicable years.

See [Notes for data users](#) and [States conducting surveillance, by year](#).

[http://apps.nccd.cdc.gov/brfss/age\\_c.asp?qttype=C&qttype\\_c=C&bkey=20045308&yr=2004&state=WA&q...](http://apps.nccd.cdc.gov/brfss/age_c.asp?qttype=C&qttype_c=C&bkey=20045308&yr=2004&state=WA&q...) 3/6/2006

TX315\_007

## At least one drink of alcohol in past 30 days Washington vs Nationwide - 2004 Age: 18-24



Compare Washington - 2004 to Another Year:

2004

Compare

Compare Washington - 2004 to Another State:

Washington

Compare

[Privacy Policy](#) | [Accessibility](#)[BRFSS Home](#) | [Contact Us](#)[CDC Home](#) | [Search](#) | [Health Topics A-Z](#)

This page last reviewed June 15, 2005

[United States Department of Health and Human Services](#)  
[Centers for Disease Control and Prevention](#)  
[National Center Chronic Disease Prevention and Health Promotion](#)  
[Division of Adult and Community Health](#)